

Ontogenetic effects on stable carbon and oxygen isotopes in tests of live (Rose Bengal stained) benthic foraminifera from the Pakistan continental margin

Submitted by Emmanuel Lemoine on Tue, 09/16/2014 - 11:50

Titre	Ontogenetic effects on stable carbon and oxygen isotopes in tests of live (Rose Bengal stained) benthic foraminifera from the Pakistan continental margin
Type de publication	Article de revue
Auteur	Schumacher, Stefanie [1], Jorissen, Frans [2], Mackensen, Andreas [3], Gooday, Andrew J. [4], Pays, Olivier [5]
Editeur	Elsevier
Type	Article scientifique dans une revue à comité de lecture
Année	2010
Langue	Anglais
Date	2010/09
Numéro	3-4
Pagination	92 - 103
Volume	76
Titre de la revue	Marine Micropaleontology
ISSN	0377-8398
Mots-clés	Arabian Sea [6], Live benthic foraminifera [7], Ontogenetic effect [8], Oxygen and carbon stable isotopes [9], Oxygen minimum zone [10], Test size [11]
Résumé en anglais	<p>We determined the stable oxygen and carbon isotopic composition of live (Rose Bengal stained) tests belonging to different size classes of two benthic foraminiferal species from the Pakistan continental margin. Samples were taken at two sites, with water depths of about 135 and 275 m, corresponding to the upper boundary and upper part of the core region of the oxygen minimum zone (OMZ). For <i>Uvigerina</i> ex gr. <i>Uvigerina semiornata</i> and <i>Bolivina</i> aff. <i>Bolivina dilatata</i>, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values increased significantly with increasing test size. In the case of <i>Uvigerina</i> ex gr. <i>U. semiornata</i>, $\delta^{13}\text{C}$ increased linearly by about 0.105‰ for each 100-μm increment in test size, whereas $\delta^{18}\text{O}$ increased by 0.02 to 0.06‰ per 100 μm increment. For <i>Bolivina</i> aff. <i>B. dilatata</i> the relationship between test size and stable isotopic composition is better described by logarithmic equations. A strong positive linear correlation is observed between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of both taxa, with a constant ratio of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values close to 2:1. This suggests that the strong ontogenetic effect is mainly caused by kinetic isotope fractionation during CO_2 uptake. Our data underline the necessity to base longer $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ isotope records derived from benthic foraminifera on size windows of 100 μm or less. This is already common practice in down-core isotopic studies of planktonic foraminifera.</p>
URL de la notice	http://okina.univ-angers.fr/publications/ua3902 [12]
DOI	10.1016/j.marmicro.2010.06.002 [13]

Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=6621](http://okina.univ-angers.fr/publications?f[author]=6621)
- [2] <http://okina.univ-angers.fr/f.jorissen/publications>
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=6622](http://okina.univ-angers.fr/publications?f[author]=6622)
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=6408](http://okina.univ-angers.fr/publications?f[author]=6408)
- [5] <http://okina.univ-angers.fr/olivier.pays/publications>
- [6] [http://okina.univ-angers.fr/publications?f\[keyword\]=8127](http://okina.univ-angers.fr/publications?f[keyword]=8127)
- [7] [http://okina.univ-angers.fr/publications?f\[keyword\]=8109](http://okina.univ-angers.fr/publications?f[keyword]=8109)
- [8] [http://okina.univ-angers.fr/publications?f\[keyword\]=8229](http://okina.univ-angers.fr/publications?f[keyword]=8229)
- [9] [http://okina.univ-angers.fr/publications?f\[keyword\]=8230](http://okina.univ-angers.fr/publications?f[keyword]=8230)
- [10] [http://okina.univ-angers.fr/publications?f\[keyword\]=8131](http://okina.univ-angers.fr/publications?f[keyword]=8131)
- [11] [http://okina.univ-angers.fr/publications?f\[keyword\]=8231](http://okina.univ-angers.fr/publications?f[keyword]=8231)
- [12] <http://okina.univ-angers.fr/publications/ua3902>
- [13] <http://dx.doi.org/10.1016/j.marmicro.2010.06.002>

Publié sur *Okina* (<http://okina.univ-angers.fr>)